IFT-UAM/CSIC-08-19

arXiv:0804.0336v2 [hep-ph] 22 Dec 2008

Sterile neutrino dark matter in $B-L$ extension of the standard model and galactic 511 keV line

Shaaban Khalil*
Centre for Theoretical Physics, The British University in Egypt,
El Sherouk City, Postal No. 11837, P.O. Box 43, Egypt. and
Department of Mathematics, Ain Shams University,
Faculty of Science, Cairo, 11566, Egypt.

Osamu Seto †‡
Instituto de Física Teórica UAM/CSIC,
Universidad Autónoma de Madrid, Cantoblanco, Madrid 28049, Spain.

(Dated: December 22, 2008)

Abstract

Sterile right-handed neutrinos can be naturally embedded in a low scale gauged $U(1)_{B-L}$ extension of the standard model. We show that, within a low reheating scenario, such a neutrino is an interesting dark matter candidate. We emphasize that if the neutrino mass is of the order of MeV, then it accounts for the measured dark matter relic density and also accommodates the observed flux of 511 keV photons from the galactic bulge.

PACS numbers: 95.35.+d, 13.15.+g, 98.80.Cq, 98.70.Rz

[†] Present address: William I. Fine Theoretical Physics Institute, University of Minnesota, Minneapolis, MN 55455, USA

[*] Electronic address: skhalil@bue.edu.eg

[‡] Electronic address: seto@physics.umn.edu

I. INTRODUCTION

The existence of non-baryonic dark matter (DM) and non-vanishing neutrino masses [1], based on the observation of neutrino oscillation, are considered as the most important evidences for new physics beyond the standard model (SM). Here, we argue that these two exciting problems may be simultaneously solved in a simple extension of the SM, that is based on the gauge group: $G_{B-L} = SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$ [2, 3]. In this class of models, three SM singlet fermions arise naturally due to the $U(1)_{B-L}$ anomaly cancellation condition. These singlet fermions are regarded as the right-handed neutrinos [3]. The scale of the right-handed neutrino masses is no longer arbitrary, it is proportional to the scale of $B-L$ symmetry breaking. It was shown that, similarly to the electroweak symmetry, the scale of $B-L$ can be linked with the supersymmetry breaking scale at the observed sector and then the $B-L$ symmetry is radiatively broken at TeV scale [4].

This model of low scale $B-L$ extension of the SM can account for the experimental results for the small neutrino masses and their large mixing through a low scale seesaw mechanism [5] and Dirac neutrino Yukawa couplings of order $\lesssim 10^{-6}$ [6]. Moreover, this model predicts an extra neutral gauge boson Z' corresponding to the $B-L$ gauge symmetry and an extra SM singlet scalar (extra Higgs). The phenomenology of this model and its potential discovery at the Large Hadron Collider (LHC) has been studied [7, 8].

As emphasized in [3], a right-handed neutrino ν_R has tree level interactions with the SM leptons L, SM Higgs ϕ, Z', and extra Higgs χ. They are described by

$$\mathcal{L}^{\rm int}_{\nu_R} = q_{\nu_R} g_{B-L} \bar{\nu}_R Z'_\mu \sigma^\mu \nu_R - (\lambda_\nu \bar{L} \tilde{\phi} \nu_R + \frac{1}{2} \lambda_{\nu_R} \bar{\nu}^c_R \chi \nu_R + \text{h.c.}). \tag{1}$$

The first term is due to the covariant derivative for right-handed neutrinos, g_{B-L} is the $B-L$ gauge coupling and q_{ν_R} is its charge. Other interaction terms refer to the possible Yukawa interactions that involve the right-handed neutrinos. After the symmetry breakdown, the right-handed neutrinos acquire Majorana masses

$$M_{\nu_{R_i}} = \lambda_{\nu_{R_i}} \langle \chi \rangle \equiv \frac{1}{\sqrt{2}} \lambda_{\nu_{R_i}} v'. \tag{2}$$

For v' of $\mathcal{O}(1)$ TeV, the right-handed neutrino masses may vary, according to the values of the Yukawa couplings $\lambda_{\nu_{R_i}}$, from TeV scale to a lighter scale. If the right-handed neutrinos are subjected to the flavour symmetry that should account for the quark and lepton mass hierarchy, then it is very natural to find $M_{\nu_{R_1}} \ll M_{\nu_{R_2}} < M_{\nu_{R_3}}$. In this case, it becomes quite plausible that the mass of the lightest right-handed neutrino is of order of MeV [1].

In this paper, we consider the scenario where a right-handed neutrino can be a DM candidate with the sufficient abundance. In addition, it provides a natural explanation for the flux of 511 keV photons from the galactic bulge observed by INTEGRAL satellite [10]. It was suggested that this emission might be originated from the annihilation [11, 12, 13, 14, 15] or decay [16, 17, 18, 19, 20, 21, 22, 23] of DM particles into low energy electron-positron pairs. The outgoing positron loses its kinetic energy by collisions with baryonic materials and eventually forms positronium with an electron in the medium. This positronium would annihilate into monoenergetic photons of 511 keV. Therefore, it was concluded that the DM candidates have to be quite light, perhaps as light as electron; otherwise the positrons produced are too energetic to form positronium. Conservative constraints imply that the mass of the corresponding DM should be of $\mathcal{O}(10)$ MeV.

The paper is organized as follows. In section 2 we show that the lightest right-handed neutrinos can be produced in enough quantities to be the main component of dark matter in a thermal history with order MeV reheating temperature. In section 3 we discuss the possibility that the MeV right-handed neutrino in $B-L$ extension of the SM can be a source for the observed 511 keV gamma ray line. Finally we give our conclusion and summary remarks in section 4.

II. DARK MATTER

In this section we consider the lightest right-handed neutrino in TeV scale $B-L$ extension of the SM as a DM candidate.

First, one should notice that this dark matter candidate is not stable. This is because, to be precise, the dark matter candidate is the mixture of a left-handed neutrino ν_L and a right-handed neutrino ν_R in the mass eigen states obtained by diagonalizing the mass matrix

$$M(\nu_L, \nu_R) = \begin{pmatrix} 0 & m_D^T \\ m_D & M_{\nu_R} \end{pmatrix}, \tag{3}$$

where m_D is the Dirac neutrino mass term. Although the lightest right-handed-like neutrino N_1 is a candidate of dark matter, it contains a tiny component of a left-handed neutrino that interacts through the weak gauge bosons $W^\pm$ and Z. Thus, even if N_1 is ligher than the SM-like Higgs, it may still decay into lepton, anti-lepton and neutrino via Z and W exchange. We will examine this decay in detail in the Sec. 4.

---

[1] The smallness might be a consequence of an extra dimension [9]

Hence, the sterile (right-handed) neutrino can be considered as DM candidate if it is rather light, around MeV or less, and if the mixing with light neutrinos is very tiny, in order to have a sufficiently long lifetime. Furthermore, the radiative decay $N_1 \to \nu + \gamma$ often gives further stringent constraints, for keV $-$ MeV sterile neutrinos, on the left-right mixing due to the line γ-ray and x-ray background radiation [24, 25]. Therefore, from now on, we assume that the DM candidate right-handed neutrino has a small enough mixing with left-handed neutrino to satisfy these constraints.

It has been assumed that sterile neutrino dark matter production is realized by so-called Dodelson-Widrow (DW) mechanism [26] due to the mixing oscillation between active and sterile neutrinos [26, 27]. However, the recent stringent bound from x-ray background seems to exclude such unstable sterile neutrino dark matter with the minimal production scenario via DW mechanism [28, 29]. To be precise, for a small enough active-sterile mixing to be consistent with the x-ray background constraints, the abundance of sterile neutrino produced by DW mechanism is too low to account for whole dark matter [30]. Hence, another production mechanism is necessary if one wants to construct a sterile neutrino as the principal component of dark matter. For instance, Shaposhnikov and Tkachev [31], and Kusenko [32, 33] proposed the production of sterile neutrinos by the decay of new gauge singlet scalar particle. This is also one fact leading us to introduce a new interaction to produce sterile neutrinos.

It is remarkable that N_1 in this model can not be thermal relics. N_1 has the gauge interaction of $U(1)_{B-L}$ with apparently heavy Z' exchange [2]. Then, the thermal relic abundance is qualitatively same as that of heavy left-handed neutrinos. Here, Z' essentially plays the role of the weak boson in the standard [34, 35]. If such a light N_1 is thermal relic, it becomes hot dark matter. Therefore, we will consider a sort of non-thermal production under a non-standard thermal history with a low reheating temperature T_R. Such a thermal history may be realized by some dominated late decaying object such as a moduli field or a late mini inflation [36, 37]. The studies of sterile neutrino dark matter under such a thermal history were also done [38, 39, 40].

The leading diagram for N production is illustrated in figure 1. As can be seen from the diagram, in $B-L$ extension of the SM, the production of MeV right-handed neutrino is dominated by the Z' exchange contribution from leptons and left-handed neutrinos. The effects of intermediate H' and H scalar Higgs bosons are strongly suppressed and can be safely neglected.

---

[2] Z' can be very light, if g_{B-L} is also very small.



FIG. 1: Feynman diagrams in $B-L$ extension of the SM that dominantly contribute to the right-handed neutrino production by the Z' exchange.

In the production, a small ν_L component in N_1 is not important at all, since the oscillation production is not efficient in our scenario. Hence, in the rest of this section, we consider $N_1 = N_R$, where N_R denotes a right-handed neutrino in the four component Majorana representation $N_R = \nu_R + \nu_R^c$ which satisfies $N_R^c = N_R$.

The relic abundance can be estimated by solving the Boltzmann equation

$$\dot{n} + 3Hn = \langle \sigma v(ij \to NN) \rangle n_i n_j . \tag{4}$$

n is the number density of N. $\langle \sigma v \rangle$ is the thermally averaged product of the pair production cross section of N and the relative velocity with the initial state particles i and j. For the calculation of the production cross section of right-handed neutrinos from the initial state i and j particles, we introduce $w(s)$ defined by

$$w(s; ij \to) \equiv \frac{1}{32\pi} \int \frac{d\cos\theta}{2} \sqrt{\frac{s - 4m_{\rm in}^2}{s}} |\bar{\mathcal{M}}_{ij}(s, \cos\theta)|^2 , \tag{5}$$

where s is the usual Mandelstam variable, θ is the scattering angle in the center of mass frame, $m_{\rm in}$ is the mass of the initial state particle, and $|\bar{\mathcal{M}}|^2$ is the squared amplitude matrix after the spin summing and averaging and also some multiplications for identical particles. Then, the right-hand side of the Boltzmann equation can be rewritten as

$$\langle \sigma v(ij \to NN) \rangle n_i n_j = \frac{4T^4}{(2\pi)^5} \int_{\frac{2m_N}{T}}^{\infty} dx\, x^2 K_1(x) \sum_{(i,j)} g_i g_j w(x^2 T^2; ij \to) , \tag{6}$$

with $x \equiv \sqrt{s}/T$ and $K_1(x)$ being the first order modified Bessel function. Here g_i stand for the internal degrees of freedoms of the i particle.

As we will show, the suitable reheating temperature is about several MeV. Hence, the relevant initial states for the production are $\nu\,\bar{\nu}$ and e^+e^-. For each initial state, we found

$$\int \frac{d\cos\theta}{2}|\bar{\mathcal{M}}_{\nu\bar{\nu}}(s,\cos\theta)|^2 = \frac{2}{3}\frac{|g_{B-L}^2 q_N q_\nu|^2}{(s-M_{Z'}^2)^2 + M_{Z'}^2\Gamma_{Z'}^2}(s-4m_N^2)s, \tag{7}$$

$$\int \frac{d\cos\theta}{2}|\bar{\mathcal{M}}_{e^+e^-}(s,\cos\theta)|^2 = \frac{4}{3}\frac{|g_{B-L}^2 q_N q_e|^2}{(s-M_{Z'}^2)^2 + M_{Z'}^2\Gamma_{Z'}^2}(s-4m_N^2)(s+2m_e^2), \tag{8}$$

where $q_N = q_e = q_\nu = -1$ are the $B-L$ charges respectively. The gauge boson Z' has the mass $M_{Z'}$, would be of order $\mathcal{O}(100)$ GeV for $g_{B-L} = \mathcal{O}(0.1)$, and its decay width $\Gamma_{Z'}$.

By introducing $Y \equiv n/s$ with s being the entropy density and using the relation $\dot{T} = -HT$, we found the final abundance as

$$\begin{aligned} Y_\infty &= \int^{T_R} dT \frac{\langle\sigma v(ij\to NN)\rangle n_i n_j}{sHT} dT \\ &\simeq 3.5\times 10^{-7}\left(\frac{10}{g_*}\right)^{3/2}\left(\frac{T_R}{5\,\mathrm{MeV}}\right)^3\left(\frac{9.7\,\mathrm{TeV}}{M_{Z'}/g_{B-L}}\right)^4, \end{aligned} \tag{9}$$

for our range of interest $M_{Z'} \gg T_R \gg m_N$. This can be rewritten in terms of the density parameter as

$$\Omega_N h^2 \simeq 0.1 \times \left(\frac{m_N}{1\,\mathrm{MeV}}\right)\left(\frac{Y_\infty}{3.5\times 10^{-7}}\right). \tag{10}$$

One can see from this equation that the heavier N, the larger $\Omega_N h^2$ obtained. For given T_R and $M_{Z'}/g_{B-L}$, if the mass of N_i is larger than the value corresponding to $\Omega_N h^2 = 0.1$ and lower than T_R [3], then such heavier N_i has to be unstable and decay in order not to behave as dark matter.

For $m_N \simeq 1$ MeV, the experimental lower bound $M_{Z'}/g_{B-L} \gtrsim 6$ TeV corresponds to the lower bound on the reheating temperature $T_R \gtrsim 2.7$ MeV, form equation (9). Since the lower bound is about 2 MeV [41], the consistency with the big bang nucleosynthesis is ensured for $m_N \sim 1$ MeV.

Finally, one may notice that the neutrinos produced have a thermal spectrum though they are not thermal relic, in other words, they were not in equilibrium, because the initial state particles in the scattering processes of the production, neutrinos and e^+, e^-, are in thermal equilibrium. This is simply same as the thermal production of gravitino dark matter by scatterings in a thermal bath [42]. Hence, if the right-handed neutrinos are light enough, they could be warm dark matter and the mass would be constrained as $m_N \gtrsim 10$ keV by the Lyman alpha, as usual.

---

[3] Needless to say, for $m_{N_i} > T_R$, such a heavy neutrino production is kinematically blocked.

III. INTEGRAL ANOMALY

We now consider the possibility that our right-handed neutrino with $\mathcal{O}$(MeV) mass in $B-L$ extension of the SM can account for the observed flux of 511 keV photons. As mentioned above, the SPI spectrometer on the INTEGRAL satellite has detected an intense 511 keV gamma ray line flux which, at 2σ level, is given by [10]:

$$\Phi_{\gamma,511} = (1.05 \pm 0.06) \times 10^{-3} \text{photon cm}^{-2} s^{-1}, \tag{11}$$

with a width of 3 keV.

The flux of the gamma ray may be explained through the decay of right-handed-like neutrino dark matter. The interaction is given by

$$\mathcal{L}_{\rm int} = \bar{\nu}_i (U_{\alpha i})^\dagger U_{\alpha I} \gamma^\mu \left(\frac{g}{2\cos\theta_W} Z_\mu + g_{B-L} q_\nu Z'_\mu \right) N_I + \bar{l^-}_\alpha \gamma^\mu \frac{g}{\sqrt{2}} W^- U_{\alpha I} N_I + \text{h.c.}, \tag{12}$$

where ν_i and N_I are left- and right-handed-like neutrino in the mass eigen state respectively. Both i and I run over $1,2,3$. θ_W is the Weinberg angle, and U is the unitary matrix for transforming neutrinos from the flavour and Majorana mass eigenstate into the actual mass eigenstate with α being left-handed flavour and right-handed Majorana mass index, in other words, α runs $e, \mu, \tau, R_1, R_2, R_3$. The rate of decay of dark matter N_1 into e^+, e^-, ν is estimated as

$$\begin{aligned} \Gamma(N_1 \to e, \bar{e}, \nu) \ \simeq \ & \frac{G_F^2 m_N^5}{192\pi^3} \sum_i \left| \sum_\alpha (U_{\alpha i})^\dagger U_{\alpha 1} \right|^2 \\ & \times \left[\frac{c_A^2}{2} + \frac{(-c_A)}{2} \left(\frac{(U_{ei})^\dagger U_{eN_1}}{\sum_\alpha (U_{\alpha i})^\dagger U_{\alpha N_1}} + \text{c.c.} \right) + \left| \frac{(U_{ei})^\dagger U_{eN_1}}{\sum_\alpha (U_{\alpha i})^\dagger U_{\alpha N_1}} \right|^2 \right], \end{aligned} \tag{13}$$

in the $m_e \ll m_N$ limit, with G_F being the Fermi constant. Here, the Z' gauge boson exchange contribution is omitted because it is much smaller than the others. In addition, the vector contribution of the neutral current is neglected because $(c_V, c_A) = (\mathcal{O}(0.01), -1/2)$ for the electron. We can regard the factor $\sum_i \left| \sum_\alpha (U_{\alpha i})^\dagger U_{\alpha 1} \right|^2$ as $\sin^2 2\theta/4$ in the standard notation for the mixing of a sterile neutrino.

It has been shown that decaying dark matter produces the 511 keV gamma ray flux of [16, 17]

$$\Phi_{511\gamma} \sim 10^{-3} \left(\frac{10^{27}\,\text{s}}{\Gamma(N_1 \to e\bar{e}\nu)^{-1}} \right) \left(\frac{1\,\text{MeV}}{m_N} \right) \text{cm}^{-2} s^{-1}. \tag{14}$$

From equations (13) and (14) with $((U_{ei})^\dagger U_{eN_1})/(\sum_\alpha (U_{\alpha i})^\dagger U_{\alpha N_1}) \leq 1$, one finds that the observed 511 keV line gamma ray can be explained if the mixing angle is of order:

$$\sin^2 2\theta \left(\frac{m_N}{1\text{MeV}} \right)^4 \simeq 10^{-22}, \tag{15}$$

which is consistent with the small mixing between light and heavy neutrinos in such models.

IV. SUMMARY

We pointed out a new mechanism of production of sterile neutrino by new $U(1)_{B-L}$ gauge boson Z' under a kind of non-standard thermal history with a low reheating temperature. If the reheating temperature is low enough, even if there is an additional gauge interaction, sterile neutrinos cannot reach thermal equilibrium. Hence, these are produced only through the scattering in the thermal bath. By adjusting the reheating temperature T_R and the scale of $B-L$, $M_{Z'}/g_{B-L}$, sufficient amounts of sterile neutrinos can be produced for rather wide range of mass.

In addition, we also have shown it can account for 511 keV gamma ray from galactic bulge claimed by INTEGRAL/SPI, for $m_N \gtrsim 1$ MeV. Picciotto and Pospelov have also explained this via decaying sterile neutrino. However, in their model, sterile neutrino cannot be abundant enough to account for whole dark matter energy density since they assumed a DW mechanism for the production [17]. This is because the left-right mixing angle is responsible for both the production and the decay of dark matter ν_R. For given $\sin^2 2\theta$ suitable for sufficient production of dark matter via the DW mechanism, the predicted flux turned out too strong. Thus, another additional dark matter component is needed to make up for the lack of the dark matter energy density. Some other models also require additional species of dark matter, in other words multi-component dark matter, to obtain $\Omega_{DM}h^2 \simeq 0.1$ [18, 20] for a similar reason, namely too strong flux. In contrast, in our scenario, a single component of dark matter is enough to explain both the cosmological abundance and the flux of 511 keV line gamma ray. On the other hand, the price to pay is to assume a low reheating temperature scenario, where successful baryogenesis is hard in general, though Affleck-Dine mechanism [43] might be applicable [44] if we supersymmetrize the model as in [4].

Acknowledgments

OS is grateful for warm hospitality at Centre for Theoretical Physics in The British University in Egypt where this work was initiated. The work of OS is supported by the MEC project FPA 2004-02015, the Comunidad de Madrid project HEPHACOS (No. P-ESP-00346). The work of SK was partially supported by the ICTP grant project 30 and the Egyptian Academy of scientific

research and technology.

---

[1] For a review, see e.g., R. N. Mohapatra *et al.*, Rept. Prog. Phys. **70**, 1757 (2007).

[2] R. N. Mohapatra and R. E. Marshak, Phys. Rev. Lett. **44**, 1316 (1980) [Erratum-ibid. **44**, 1643 (1980)]; R. E. Marshak and R. N. Mohapatra, Phys. Lett. B **91**, 222 (1980); C. Wetterich, Nucl. Phys. B **187**, 343 (1981).

[3] S. Khalil, J. Phys. G. **35** 055001 (2008).

[4] S. Khalil and A. Masiero, Phys. Lett. B **665**, 374 (2008).

[5] T. Yanagida, in *Proceedings of Workshop on the Unified Theory and the Baryon Number in the Universe*, Tsukuba, Japan, edited by A. Sawada and A. Sugamoto (KEK, Tsukuba, 1979), p 95; M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, Proceedings of Workshop, Stony Brook, New York, 1979, edited by P. Van Nieuwenhuizen and D. Z. Freedman (North-Holland, Amsterdam, 1979), p 315; R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).

[6] M. Abbas and S. Khalil, JHEP **0804**, 056 (2008).

[7] W. Emam and S. Khalil, Eur. Phys. J. C **522**, 625 (2007).

[8] K. Huitu, S. Khalil, H. Okada and S. K. Rai, arXiv:0803.2799 [hep-ph].

[9] K. Kadota, Phys. Rev. D **77**, 063509 (2008).

[10] J. Knodlseder *et al.*, Astron. Astrophys. **411**, L457 (2003); P. Jean *et al.*, Astron. Astrophys. **407**, L55 (2003).

[11] C. Boehm, D. Hooper, J. Silk, M. Casse and J. Paul, Phys. Rev. Lett. **92**, 101301 (2004).

[12] P. Fayet, Phys. Rev. D **70**, 023514 (2004).

[13] J. F. Gunion, D. Hooper and B. McElrath, Phys. Rev. D **73**, 015011 (2006).

[14] C. Boehm, J. Orloff and P. Salati, Phys. Lett. B **641**, 247 (2006).

[15] D. Hooper and K. M. Zurek, Phys. Rev. D **77**, 087302 (2008).

[16] D. Hooper and L. T. Wang, Phys. Rev. D **70**, 063506 (2004).

[17] C. Picciotto and M. Pospelov, Phys. Lett. B **605**, 15 (2005).

[18] M. Kawasaki and T. Yanagida, Phys. Lett. B **624**, 162 (2005).

[19] F. Takahashi and T. T. Yanagida, Phys. Lett. B **635**, 57 (2006).

[20] S. Kasuya and F. Takahashi, Phys. Rev. D **72**, 085015 (2005).

[21] S. Kasuya and M. Kawasaki, Phys. Rev. D **73**, 063007 (2006).

[22] E. J. Chun and H. B. Kim, JHEP **0610**, 082 (2006).

[23] M. Pospelov and A. Ritz, Phys. Lett. B **651**, 208 (2007).

[24] A. Boyarsky, D. Iakubovskyi, O. Ruchayskiy and V. Savchenko, Mon. Not. Roy. Astron. Soc. **387**, 1361 (2008).

[25] A. Boyarsky, D. Malyshev, A. Neronov and O. Ruchayskiy, Mon. Not. Roy. Astron. Soc. **387**, 1345 (2008).

[26] S. Dodelson and L. M. Widrow, Phys. Rev. Lett. **72**, 17 (1994).

[27] A. D. Dolgov and S. H. Hansen, Astropart. Phys. **16**, 339 (2002).

[28] T. Asaka, M. Laine and M. Shaposhnikov, JHEP **0606**, 053 (2006) and JHEP **0701**, 091 (2007).

[29] H. Yuksel, J. F. Beacom and C. R. Watson, Phys. Rev. Lett. **101** 121301 (2008).

[30] A. Palazzo, D. Cumberbatch, A. Slosar and J. Silk, Phys. Rev. D **76**, 103511 (2007).

[31] M. Shaposhnikov and I. Tkachev, Phys. Lett. B **639**, 414 (2006).

[32] A. Kusenko, Phys. Rev. Lett. **97**, 241301 (2006).

[33] K. Petraki and A. Kusenko, Phys. Rev. D **77**, 065014 (2008).

[34] B. W. Lee and S. Weinberg, Phys. Rev. Lett. **39**, 165 (1977).

[35] E. W. Kolb and M. S. Turner, *The Early Universe*, Addison-Wesley (1990).

[36] G. F. Giudice, E. W. Kolb and A. Riotto, Phys. Rev. D **64**, 023508 (2001).

[37] e.g., S. Khalil, C. Munoz and E. Torrente-Lujan, New J. Phys. **4** (2002) 27; C. Pallis, Astropart. Phys. **21** (2004) 689; G. B. Gelmini and P. Gondolo, Phys. Rev. D **74**, 023510 (2006).

[38] G. Gelmini, S. Palomares-Ruiz and S. Pascoli, Phys. Rev. Lett. **93**, 081302 (2004).

[39] C. E. Yaguna, JHEP **0706**, 002 (2007).

[40] G. Gelmini, E. Osoba, S. Palomares-Ruiz and S. Pascoli, arXiv:0803.2735 [astro-ph].

[41] M. Kawasaki, K. Kohri and N. Sugiyama, Phys. Rev. Lett. **82**, 4168 (1999) and Phys. Rev. D **62**, 023506 (2000); K. Ichikawa, M. Kawasaki and F. Takahashi, Phys. Rev. D **72**, 043522 (2005).

[42] M. Bolz, W. Buchmuller and M. Plumacher, Phys. Lett. B **443**, 209 (1998); M. Bolz, A. Brandenburg and W. Buchmuller, Nucl. Phys. B **606**, 518 (2001) [Erratum-ibid. B **790**, 336 (2008)] ; F. D. Steffen, JCAP **0609**, 001 (2006).

[43] I. Affleck and M. Dine, Nucl. Phys. B **249**, 361 (1985).

[44] E. D. Stewart, M. Kawasaki and T. Yanagida, Phys. Rev. D **54**, 6032 (1996); B. A. Campbell, M. K. Gaillard, H. Murayama and K. A. Olive, Nucl. Phys. B **538**, 351 (1999); A. D. Dolgov, K. Kohri, O. Seto and J. Yokoyama, Phys. Rev. D **67**, 103515 (2003).